

Poincaré–Lelong Approach to Universality and Scaling of Correlations Between Zeros

Pavel Bleher^{1,*}, Bernard Shiffman^{2,**}, Steve Zelditch^{2,***}

¹ Department of Mathematical Sciences, IUPUI, Indianapolis, IN 46202, USA.

E-mail: bleher@math.iupui.edu

² Department of Mathematics, Johns Hopkins University, Baltimore, MD 21218, USA.

E-mail: shiffman@math.jhu.edu; zel@math.jhu.edu

Received: 17 March 1999 / Accepted: 5 August 1999

Abstract: This note is concerned with the scaling limit as $N \rightarrow \infty$ of n -point correlations between zeros of random holomorphic polynomials of degree N in m variables. More generally we study correlations between zeros of holomorphic sections of powers L^N of any positive holomorphic line bundle L over a compact Kähler manifold. Distances are rescaled so that the average density of zeros is independent of N . Our main result is that the scaling limits of the correlation functions and, more generally, of the “correlation forms” are universal, i.e. independent of the bundle L , manifold M or point on M .

Introduction

This note is a companion to our article [BSZ], in which we study the correlations between the zeros of a random holomorphic section $s \in H^0(M, L^N)$ of a power L^N of a positive line bundle $L \rightarrow M$ over a compact m -dimensional complex manifold M . Since the hypersurface volume of the zeros of a section of L^N in a ball U around a given point z_0 is $\sim N \text{Vol}(U)$, we rescale $U \rightarrow \sqrt{N}U$ to get a density of zeros independent of N . After expanding U this way, all manifolds and line bundles appear asymptotically alike, and it is natural to ask if the *local statistics* of zeros are universal, i.e. independent of L , M , ω and z_0 . To define our statistics, we first provide $H^0(M, L^N)$ with a natural Gaussian measure (see Sects. 1.1–1.2). The local statistics are measured by the scaled n -point zero correlation forms $\mathbf{K}_n^N(\frac{z^1}{\sqrt{N}}, \dots, \frac{z^n}{\sqrt{N}})$, $z^j \in \sqrt{N}U$ (see Sect. 1.3). They are smooth off the diagonal, and their norms define scaled zero correlation measures $\tilde{K}_n^N(\frac{z^1}{\sqrt{N}}, \dots, \frac{z^n}{\sqrt{N}})$. (The scaled correlation forms extend to all of $(\sqrt{N}U)^n$ as currents

* Research partially supported by NSF grant #DMS-9623214.

** Research partially supported by NSF grant #DMS-9800479.

*** Research partially supported by NSF grant #DMS-9703775.

of order 0.) The correlation measures \tilde{K}_n^N give the expected value of the product of the volumes of the zero set in n domains in M ; see Eq. (12). In [BSZ], we used geometric probability methods and a (universal) scaled Szegő kernel to prove that there exist universal limits as $N \rightarrow \infty$ of these correlation measures and more generally of the correlations between simultaneous zeros of $k \leq m$ sections. Here we use a complex analytic approach based on the Poincaré–Lelong formula for the currents of integration over the zero set of a section, together with the scaled Szegő kernel from [BSZ], to prove universality for the correlation forms. This approach, although limited to the hypersurface case, allows for a result on the level of forms and a somewhat simpler proof.

Our universality theorem is as follows:

Main Theorem. *There is a universal current $\mathbf{K}_n^\infty \in \mathcal{D}'^{(m-1)n, (m-1)n}(\mathbb{C}^{mn})$ such that the following holds: suppose that (L, h) is a positive Hermitian line bundle on an m -dimensional compact complex manifold M , and let \mathbf{K}_n^N be the n -point zero correlation current on M^n . Suppose $z^0 \in M$ and choose local holomorphic coordinates in M about z^0 such that $\Theta_h|_{z^0} = \partial\bar{\partial}|z|^2$. Then*

$$\mathbf{K}_n^N \left(\frac{z^1}{\sqrt{N}}, \dots, \frac{z^n}{\sqrt{N}} \right) = \mathbf{K}_n^\infty(z^1, \dots, z^n) + O\left(\frac{1}{\sqrt{N}}\right).$$

Furthermore, \mathbf{K}_n^∞ is a smooth form on the off-diagonal domain \mathcal{G}_n^m consisting of n -tuples of distinct points z^1, \dots, z^n in \mathbb{C}^m . The error term has k^{th} order derivatives $\leq \frac{C_{A,k}}{\sqrt{N}}$ on each compact subset $A \subset \mathcal{G}_n^m$, $\forall k \geq 0$.

This result is new in all dimensions, and the proof does not simplify in any essential way in the case $m = 1$ of Riemann surfaces. Our method leads to integral formulae for the universal limit forms, although the details rapidly become complicated as the number n of points increase. For the case $n = 2$ of the pair correlation function, we carry out the calculation in complete detail in dimension $m = 1$ and also use the method to obtain an explicit formula for the scaling limit pair correlation measures in all dimensions (Theorems 4.1 and 4.2). In particular, our formula gives the scaling limit pair correlations for $SU(m+1)$ -polynomials (which are sections of powers of the hyperplane bundle over complex projective space $\mathbb{C}\mathbb{P}^m$). The universal formula in dimension $m = 1$ agrees, as it must, with that of Hannay [Ha] in the case of random $SU(2)$ -polynomials. (The universality of this formula was not proved previously). Similar formulas for correlations of zeros of real polynomials were given in [BD].

Before we get started on the proof, the following heuristic remark on correlation measures and forms may be helpful. Although the definition of the correlation measures is formulated in terms of expected volumes of the zero set (or in the one-dimensional case, expected numbers of zeros) in domains of M , it can also be given a probabilistic interpretation: the probability that the zero divisor of a random section s simultaneously intersects balls of radius ε around z^1, \dots, z^n , respectively, is $\approx c\varepsilon^{2n} K_n^N(z^1, \dots, z^n)$ (where K_n^N is the correlation function given by $K_n^N dV_M = \tilde{K}_n^N$). The correlation form \mathbf{K}_n^N gives a more refined probability that the zero divisor of s has tangent hyperplanes close to n fixed complex hyperplanes in T_M .

A final remark on the term “universality”: in this paper it means independence from the details of the geometric setting, i.e. from the complex manifold, line bundle, metric, connection, etc. In random matrix theory (cf. [D]) it has a somewhat different meaning.

There, the “setting” consists of a fixed class of $N \times N$ matrices; what varies is the probability measure $d\nu_N$ on the space of matrices. Universality then means that the correlation functions in the scaling limit are the same for a broad class of measures $d\nu_N$. This notion of universality makes sense as well for random polynomials and sections. We could study the statistics of zeros relative to more general measures on $H^0(M, L^N)$ than Gaussian ones, e.g. measures of the form $d\nu_N(s) = e^{-\int_M (|\nabla s(z)|^2 + a|s(z)|^4 + b|s(z)|^2)} dV(z) ds$. Such measures are biased against sections with strong oscillations, and one could ask how that affects the statistics of zeros. However, we do not consider universality in the measure aspect in this paper.

1. Notation

We summarize here the notation from complex analysis that we will need in the proof. This notation is the same as in [SZ] and [BSZ], except that different normalizations for the metric and volume form are used in [SZ].

1.1. Complex geometry. We denote by $(L, h) \rightarrow M$ a holomorphic line bundle with smooth Hermitian metric h whose curvature form

$$\Theta_h = -\partial\bar{\partial} \log \|e_L\|_h^2, \tag{1}$$

is a positive $(1,1)$ -form. Here, e_L is a local non-vanishing holomorphic section of L over an open set $U \subset M$, and $\|e_L\|_h = h(e_L, e_L)^{1/2}$ is the h -norm of e_L . As in [BSZ], we give M the Hermitian metric corresponding to the Kähler form $\omega = \frac{\sqrt{-1}}{2} \Theta_h$ and the induced Riemannian volume form

$$dV_M = \frac{1}{m!} \omega^m. \tag{2}$$

We denote by $H^0(M, L^N)$ the space of holomorphic sections of $L^N = L \otimes \dots \otimes L$. The metric h induces Hermitian metrics h^N on L^N given by $\|s^{\otimes N}\|_{h^N} = \|s\|_h^N$. We give $H^0(M, L^N)$ the Hermitian inner product

$$\langle s_1, s_2 \rangle = \int_M h^N(s_1, s_2) dV_M \quad (s_1, s_2 \in H^0(M, L^N)), \tag{3}$$

and we write $|s| = \langle s, s \rangle^{1/2}$.

For a holomorphic section $s \in H^0(M, L^N)$, we let Z_s denote the current¹ of integration over the zero divisor of s :

$$(Z_s, \varphi) = \int_{Z_s} \varphi, \quad \varphi \in \mathcal{D}^{m-1, m-1}(M).$$

The Poincaré–Lelong formula (see e.g., [GH]) expresses the integration current of a holomorphic section $s = ge_L^{\otimes N}$ in the form:

$$Z_s = \frac{i}{\pi} \partial\bar{\partial} \log |g| = \frac{i}{\pi} \partial\bar{\partial} \log \|s\|_{h^N} + N\omega. \tag{4}$$

¹ Here, $\mathcal{D}^{p,q}(\Omega)$ denotes the space of compactly supported (p, q) forms on a complex manifold Ω . A current is an element of the dual space $\mathcal{D}^{p,q}(\Omega)' = \mathcal{D}^{m-p, m-q}(\Omega)$.

We also denote by $|Z_s|$ the Riemannian $(2m - 2)$ -volume along the regular points of Z_s , regarded as a measure on M :

$$(|Z_s|, \varphi) = \int_{Z_s^{\text{reg}}} \varphi d\text{Vol}_{2m-2} = \frac{1}{(m-1)!} \int_{Z_s^{\text{reg}}} \varphi \omega^{m-1}; \tag{5}$$

i.e., $|Z_s|$ is the ‘‘total variation measure’’ of the current of integration over Z_s :

$$|Z_s| = Z_s \wedge \frac{1}{(m-1)!} \omega^{m-1}. \tag{6}$$

1.2. Random sections and Gaussian measures. We now give $H^0(M, L^N)$ the complex Gaussian probability measure

$$d\mu(s) = \frac{1}{\pi^{d_N}} e^{-|c|^2} dc, \quad s = \sum_{j=1}^{d_N} c_j S_j^N, \tag{7}$$

where $\{S_j^N : 1 \leq j \leq d_N\}$ is an orthonormal basis for $H^0(M, L^N)$ and dc is $2d_N$ -dimensional Lebesgue measure. This Gaussian is characterized by the property that the $2d_N$ real variables $\Re c_j, \Im c_j$ ($j = 1, \dots, d_N$) are independent random variables with mean 0 and variance $\frac{1}{2}$; i.e.,

$$\mathbf{E} c_j = 0, \quad \mathbf{E} c_j c_k = 0, \quad \mathbf{E} c_j \bar{c}_k = \delta_{jk}.$$

Here and throughout this paper, \mathbf{E} denotes expectation: $\mathbf{E} \varphi = \int \varphi d\mu$.

We then regard the currents Z_s (resp. measures $|Z_s|$), as current-valued (resp. measure-valued) random variables on the probability space $(H^0(M, L^N), d\mu)$; i.e., for each test form (resp. function) φ , (Z_s, φ) (resp. $(|Z_s|, \varphi)$) is a complex-valued random variable.

Since the zero current Z_s is unchanged when s is multiplied by an element of \mathbb{C}^* , our results are the same if we instead regard Z_s as a random variable on the unit sphere $SH^0(M, L^N)$ with Haar probability measure. We prefer to use Gaussian measures in order to facilitate our computations.

1.3. Correlation currents and measures. The n -point correlation current of the zeros is the current on $M^n = M \times M \times \dots \times M$ (n times) given by

$$\mathbf{K}_n^N(z^1, \dots, z^n) := \mathbf{E} (Z_s(z^1) \otimes Z_s(z^2) \otimes \dots \otimes Z_s(z^n)) \tag{8}$$

in the sense that for any test form $\varphi_1(z^1) \otimes \dots \otimes \varphi_n(z^n) \in \mathcal{D}^{m-1, m-1}(M) \otimes \dots \otimes \mathcal{D}^{m-1, m-1}(M)$,

$$(\mathbf{K}_n^N(z^1, \dots, z^n), \varphi_1(z^1) \otimes \dots \otimes \varphi_n(z^n)) = \mathbf{E} [(Z_s, \varphi_1)(Z_s, \varphi_2) \cdots (Z_s, \varphi_n)]. \tag{9}$$

In a similar way we define the n -point correlation measures \tilde{K}_n^N as the ‘‘total variation measures’’ of the n -point correlation currents:

$$\tilde{K}_n^N(z^1, \dots, z^n) = \mathbf{K}_n^N(z^1, \dots, z^n) \wedge \frac{1}{(m-1)!} \omega_{z^1}^{m-1} \wedge \dots \wedge \frac{1}{(m-1)!} \omega_{z^n}^{m-1}. \tag{10}$$

By (6) and (10), we have

$$(\tilde{K}_n^N(z^1, \dots, z^n), \varphi_1(z^1) \dots \varphi_n(z^n)) = \mathbf{E}[(|Z_s|, \varphi_1)(|Z_s|, \varphi_2) \dots (|Z_s|, \varphi_n)], \quad (11)$$

where $\varphi_j \in \mathcal{C}^0(M)$. Equivalently,

$$\tilde{K}_n^N(U_1 \times \dots \times U_n) = \mathbf{E} \left[\prod_{j=1}^n \text{Vol}(Z_s \cap U_j) \right], \quad U_1, \dots, U_n \subset M. \quad (12)$$

Remark. In the case of pair correlation on a Riemann surface ($n = 2, \dim M = 1$), the correlation measures take the form

$$K_2^N(z, w) = [\Delta] \wedge (K_1^N(z) \otimes 1) + \kappa^N(z, w)\omega_z \otimes \omega_w \quad (N \gg 0),$$

where $[\Delta]$ denotes the current of integration along the diagonal $\Delta = \{(z, z)\} \subset M \times M$, and $\kappa^N \in \mathcal{C}^\infty(M \times M)$.

1.4. Szegő kernels. As in [Ze, SZ, BSZ] and elsewhere, we analyze the $N \rightarrow \infty$ limit by lifting it to a principal S^1 bundle $\pi : X \rightarrow M$. Let us recall how this goes.

We denote by L^* the dual line bundle to L , and define X as the circle bundle $X = \{\lambda \in L^* : \|\lambda\|_{h^*} = 1\}$, where h^* is the norm on L^* dual to h . We can view X as the boundary of the disc bundle $D = \{\lambda \in L^* : \rho(\lambda) > 0\}$, where $\rho(a) = 1 - \|\lambda\|_{h^*}^2$. The disc bundle D is strictly pseudoconvex in L^* , since Θ_h is positive, and hence X inherits the structure of a strictly pseudoconvex CR manifold. Associated to X is the contact form $\alpha = -i\partial\bar{\rho}|_X = i\bar{\partial}\rho|_X$. We also give X the volume form

$$dV_X = \frac{1}{m!} \alpha \wedge (d\alpha)^m = \alpha \wedge \pi^* dV_M. \quad (13)$$

The setting for our analysis of the Szegő kernel is the Hardy space $H^2(X) \subset \mathcal{L}^2(X)$ of square integrable CR functions on X , where we use the inner product

$$\langle F_1, F_2 \rangle = \frac{1}{2\pi} \int_X F_1 \bar{F}_2 dV_X, \quad F_1, F_2 \in \mathcal{L}^2(X). \quad (14)$$

We let $r_\theta x = e^{i\theta}x$ ($x \in X$) denote the S^1 action on X . The action r_θ commutes with the Cauchy-Riemann operator $\bar{\partial}_b$; hence $H^2(X) = \bigoplus_{N=0}^\infty H_N^2(X)$, where

$$H_N^2(X) = \{F \in H^2(X) : F(r_\theta x) = e^{iN\theta} F(x)\}.$$

A section s_N of L^N determines an equivariant function \hat{s}_N on X :

$$\hat{s}_N(z, \lambda) = (\lambda^{\otimes N}, s_N(z)), \quad (z, \lambda) \in X; \quad (15)$$

then $\hat{s}_N(r_\theta x) = e^{iN\theta} \hat{s}_N(x)$. The map $s \mapsto \hat{s}$ is a unitary equivalence between $H^0(M, L^{\otimes N})$ and $H_N^2(X)$.

We let $\Pi_N : \mathcal{L}^2(X) \rightarrow H_N^2(X)$ denote the orthogonal projection. The Szegő kernel $\Pi_N(x, y)$ is defined by

$$\Pi_N F(x) = \int_X \Pi_N(x, y) F(y) dV_X(y), \quad F \in \mathcal{L}^2(X). \quad (16)$$

It can be given as

$$\Pi_N(x, y) = \sum_{j=1}^{d_N} \widehat{S}_j^N(x) \overline{\widehat{S}_j^N(y)}, \tag{17}$$

where $S_1^N, \dots, S_{d_N}^N$ form an orthonormal basis of $H^0(M, L^N)$.

2. Scaling

In order that we may study the local nature of the random variable Z_s , we fix a point $z^0 \in M$ and choose a holomorphic coordinate chart $\Psi : \Omega, 0 \rightarrow U, z_0$ ($\Omega \subset \mathbb{C}^m, U \subset M$) such that

$$\Psi^* \omega_{z^0} = \frac{i}{2} \sum_{j=1}^m dz_j \wedge d\bar{z}_j \Big|_0. \tag{18}$$

For example, if L is the hyperplane section bundle $\mathcal{O}(1)$ over $\mathbb{C}P^m$ with the Fubini-Study metric h_{FS} , and $z_0 = (1 : 0 : \dots : 0)$, then the coordinate chart

$$\Psi : \mathbb{C}^m \rightarrow U = \{w \in \mathbb{C}P^m : w_0 \neq 0\}, \quad \Psi(z) = (1 : z_1 : \dots : z_m)$$

(i.e., $z_j = w_j/w_0$) satisfies (18).

To simplify notation, we identify U with Ω . For a current $T \in \mathcal{D}'^{p,q}(\Omega)$, we write

$$T\left(\frac{z}{\sqrt{N}}\right) = \left(\tau_{\sqrt{N}}\right)_* T \in \mathcal{D}'^{p,q}(\sqrt{N}\Omega), \quad (\tau_\lambda(z) = \lambda z).$$

(In particular, if $T = \sum T_{jk}(z) dz_j \wedge d\bar{z}_k$, then $T(\frac{z}{\sqrt{N}}) = \frac{1}{N} \sum T_{jk}(\frac{z}{\sqrt{N}}) dz_j \wedge d\bar{z}_k$.)

We define the *rescaled zero current* of $s \in H^0(M, L^N)$ by

$$\widehat{Z}_s^N(z) := Z_s\left(\frac{z}{\sqrt{N}}\right). \tag{19}$$

The scaled n -point correlation currents are then defined by:

$$\mathbf{E} \left(\widehat{Z}_s^N(z^1) \otimes \widehat{Z}_s^N(z^2) \otimes \dots \otimes \widehat{Z}_s^N(z^n) \right) = \mathbf{K}_n^N \left(\frac{z^1}{\sqrt{N}}, \dots, \frac{z^n}{\sqrt{N}} \right) \in \mathcal{D}'^{n,n}(M^n). \tag{20}$$

Following the approach of [SZ], we fix an orthonormal basis $\{S_j^N\}$ of $H^0(M, L^N)$ and write $S_j^N = f_j^N e_L^{\otimes N}$ over U . Any section in $H^0(M, L^N)$ may then be written as $s = \sum_{j=1}^{d_N} c_j f_j^N e_L^{\otimes N}$. To simplify the notation we let $f^N = (f_1^N, \dots, f_{d_N}^N) : U \rightarrow \mathbb{C}^{d_N}$ and we put

$$\sum_{j=1}^{d_N} c_j f_j = c \cdot f^N.$$

Hence

$$Z_s = \frac{\sqrt{-1}}{\pi} \partial \bar{\partial} \log |c \cdot f^N|, \quad \widehat{Z}_s^N = \frac{\sqrt{-1}}{\pi} \partial_z \bar{\partial}_z \log \left| c \cdot f^N \left(\frac{z}{\sqrt{N}} \right) \right|, \quad (21)$$

and therefore

$$\begin{aligned} \widehat{Z}_s(z^1) \otimes \cdots \otimes \widehat{Z}_s(z^n) &= \left(\frac{i}{\pi} \right)^n \partial_{z^1} \bar{\partial}_{z^1} \cdots \\ &\quad \partial_{z^n} \bar{\partial}_{z^n} \left[\log \left| c \cdot f^N \left(\frac{z^1}{\sqrt{N}} \right) \right| \cdots \log \left| c \cdot f^N \left(\frac{z^n}{\sqrt{N}} \right) \right| \right]. \end{aligned} \quad (22)$$

We then can write the rescaled correlation currents in the form

$$\begin{aligned} \mathbf{K}_n^N \left(\frac{z^1}{\sqrt{N}}, \dots, \frac{z^n}{\sqrt{N}} \right) &= \mathbf{E} \left(\widehat{Z}_s(z^1) \otimes \cdots \otimes \widehat{Z}_s(z^n) \right) \\ &= \left(\frac{i}{\pi} \right)^n \partial_{z^1} \bar{\partial}_{z^1} \cdots \partial_{z^n} \bar{\partial}_{z^n} \\ &\quad \int_{\mathbb{C}^{d_N}} \log \left| c \cdot f^N \left(\frac{z^1}{\sqrt{N}} \right) \right| \cdots \log \left| c \cdot f^N \left(\frac{z^n}{\sqrt{N}} \right) \right| \frac{e^{-|c|^2}}{\pi^{d_N}} dc. \end{aligned} \quad (23)$$

2.1. Scaling limit of the Szegő kernel. The asymptotics of the Szegő kernel along the diagonal were given by [Ti] and [Ze]:

$$\frac{\pi^m}{N^m} \Pi_N(x, x) = 1 + O(N^{-1}). \quad (24)$$

For our proof of the Main Theorem, we need the following lemma from [BSZ], which gives the “near-diagonal” asymptotics of the Szegő kernel.

Lemma 2.1. *Let $z^0 \in M$ and choose local coordinates $\{z^j\}$ in a neighborhood of z_0 so that $z^0 = 0$ and $\Theta_h(z_0) = \sum dz^j \wedge d\bar{z}^j$. Then*

$$\frac{\pi^m}{N^m} \Pi_N \left(\frac{z}{\sqrt{N}}, \frac{\theta}{N}; \frac{w}{\sqrt{N}}, \frac{\varphi}{N} \right) = e^{i(\theta-\varphi) + i\Im(z \cdot \bar{w}) - \frac{1}{2}|z-w|^2} + O(N^{-1/2}).$$

Here, (z, θ) denotes the point $e^{i\theta} \|e_L(z)\|_h e_L^*(z) \in X$, and similarly for (w, φ) . In (24) and Lemma 2.1, the expression $O(N^\alpha)$ means a term with k^{th} order derivatives $\leq C_k N^\alpha$, for all $k \geq 0$. Lemma 2.1 says that the Szegő kernel has a universal scaling limit. In fact, its scaling limit is the first Szegő kernel of the reduced Heisenberg group; see [BSZ].

3. Universality

All the ideas of the proof of the Main Theorem occur in the simplest case $n = 2$. So first we prove universality in that case and then extend the proof to general n .

Thus, our first object is to prove universality of the large N limit of the rescaled pair correlation current (from (23) with $n = 2$)

$$\begin{aligned} \mathbf{K}_2^N \left(\frac{z}{\sqrt{N}}, \frac{w}{\sqrt{N}} \right) &= \mathbf{E} \left(\widehat{Z}_s^N(z) \otimes \widehat{Z}_s^N(w) \right) \\ &= \frac{-1}{\pi^2} \partial_z \bar{\partial}_z \partial_w \bar{\partial}_w \int_{\mathbb{C}^{d_N}} \log \left| c \cdot f^N \left(\frac{z}{\sqrt{N}} \right) \right| \log \left| c \cdot f^N \left(\frac{w}{\sqrt{N}} \right) \right| \frac{e^{-|c|^2}}{\pi^{d_N}} dc. \end{aligned} \tag{25}$$

As in [SZ], we write $f^N = |f^N|u^N$ and expand the integrand in (25):

$$\begin{aligned} \log \left| c \cdot f^N \left(\frac{z}{\sqrt{N}} \right) \right| \log \left| c \cdot f^N \left(\frac{w}{\sqrt{N}} \right) \right| &= \log |f^N \left(\frac{z}{\sqrt{N}} \right)| \log |f^N \left(\frac{w}{\sqrt{N}} \right)| \\ &\quad + \log |f^N \left(\frac{z}{\sqrt{N}} \right)| \log |c \cdot u^N \left(\frac{w}{\sqrt{N}} \right)| \\ &\quad + \log |f^N \left(\frac{w}{\sqrt{N}} \right)| \log |c \cdot u^N \left(\frac{z}{\sqrt{N}} \right)| \\ &\quad + \log |c \cdot u^N \left(\frac{z}{\sqrt{N}} \right)| \log |c \cdot u^N \left(\frac{w}{\sqrt{N}} \right)|. \end{aligned} \tag{26}$$

Let us denote the terms resulting from this expansion by E_1, E_2, E_3, E_4 , respectively. In particular,

$$E_1 = \frac{-1}{\pi^2} \partial_z \bar{\partial}_z \partial_w \bar{\partial}_w \left[\log |f^N \left(\frac{z}{\sqrt{N}} \right)| \log |f^N \left(\frac{w}{\sqrt{N}} \right)| \right]. \tag{27}$$

By (15), $\widehat{S}_j^N(z, \theta) = e^{iN\theta} \|e_L(z)\|_h^N f_j^N(z)$, where (z, θ) are the coordinates in X given in Sect. 2.1. By (17),

$$\Pi_N(z, w) = \|e_L(z)\|_h^N \|e_L(w)\|_h^N \langle f^N(z), f^N(w) \rangle, \tag{28}$$

where we write $\Pi_N(z, w) = \Pi_N(z, 0; w, 0)$. Since $\Pi_N(z, z)^{1/2} = \|e_L(z)\|_h^N |f^N(z)|$, each factor in (27) has the form $\frac{1}{2} \log \Pi_N \left(\frac{z}{\sqrt{N}}, \frac{z}{\sqrt{N}} \right) - N \log \|e_L \left(\frac{z}{\sqrt{N}} \right)\|_h$. By (24),

$$\partial_z \bar{\partial}_z \partial_w \bar{\partial}_w \log \Pi_N \left(\frac{z}{\sqrt{N}}, \frac{z}{\sqrt{N}} \right) \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

On the other hand

$$-iN \partial_z \bar{\partial}_z \log \|e_L \left(\frac{z}{\sqrt{N}} \right)\|_h = \omega \left(\frac{z}{\sqrt{N}} \right).$$

Hence the first term converges to the normalized Euclidean (double) Kähler form:

$$E_1 = \frac{i}{2\pi} \partial \bar{\partial} |z|^2 \wedge \frac{i}{2\pi} \partial \bar{\partial} |w|^2 + O\left(\frac{1}{N}\right). \tag{29}$$

The middle two terms vanish since the integrals in E_2 and E_3 are independent of w and z respectively (see [SZ, §3.2]). The “interesting term” is therefore

$$E_4 = \frac{-1}{\pi^2} \partial_z \bar{\partial}_z \partial_w \bar{\partial}_w \int_{\mathbb{C}^{d_N}} \log |c \cdot u^N \left(\frac{z}{\sqrt{N}} \right)| \log |c \cdot u^N \left(\frac{w}{\sqrt{N}} \right)| \frac{e^{-|c|^2}}{\pi^{d_N}} dc. \tag{30}$$

To evaluate E_4 , we consider the integral

$$G_2^N(x^1, x^2) := \int_{\mathbb{C}^{d_N}} \log |c \cdot x^1| \log |c \cdot x^2| \frac{e^{-|c|^2}}{\pi^{d_N}} dc \quad (x^1, x^2 \in \mathbb{C}^{d_N}) \quad (31)$$

with $x^1 = u^N(\frac{z}{\sqrt{N}})$, $x^2 = u^N(\frac{w}{\sqrt{N}})$. To simplify it, we construct a Hermitian orthonormal basis $\{e_1, \dots, e_{d_N}\}$ for \mathbb{C}^{d_N} such that $x^1 = e_1$ and

$$x^2 = \xi_1 e_1 + \xi_2 e_2, \quad \xi_1 = \langle x^2, x^1 \rangle, \quad \xi_2 = \sqrt{1 - |\xi_1|^2}. \quad (32)$$

This is possible because we can always multiply e_2 by a phase $e^{i\theta}$ so that ξ_2 is positive real. We then make a unitary change of variables to express the integral in the $\{e_j\}$ coordinates. Since the Gaussian is $U(d_N)$ -invariant, (31) simplifies to

$$G_2^N(x^1, x^2) = G_2(\xi_1, \xi_2) = \frac{1}{\pi^2} \int_{\mathbb{C}^2} e^{-(|c_1|^2 + |c_2|^2)} \log |\xi_1| \log |c_1 \xi_1 + c_2 \xi_2| dc_1 dc_2 \quad (33)$$

(where we used the fact that the Gaussian integral in each c_j , $j \geq 3$, equals 1 by construction). By performing a rotation of the c_1 variable, we may replace ξ_1 with $|\xi_1|$ and replace $G_2(\xi_1, \xi_2)$ with

$$G(\cos \theta) := G_2(\cos \theta, \sin \theta), \quad (34)$$

where $\cos \theta = |\xi_1| = |\langle x^1, x^2 \rangle|$, $0 \leq \theta \leq \pi/2$. Hence (30) becomes

$$E_4 = \frac{-1}{\pi^2} \partial_z \bar{\partial}_z \partial_w \bar{\partial}_w G(\cos \theta_N), \quad \cos \theta_N = \left| \left\langle u^N\left(\frac{z}{\sqrt{N}}\right), u^N\left(\frac{w}{\sqrt{N}}\right) \right\rangle \right|. \quad (35)$$

By the universal scaling formula for the Szegő kernel (Lemma 2.1) and (28), we have

$$\cos \theta_N = \frac{|\Pi_N(z, w)|}{\Pi_N(z, z)^{1/2} \Pi_N(w, w)^{1/2}} = e^{-\frac{1}{2}|z-w|^2} + O(N^{-\frac{1}{2}}). \quad (36)$$

Thus we get the universal formula:

$$\mathbf{K}_2^\infty(z, w) = \frac{i}{2\pi} \partial \bar{\partial} |z|^2 \wedge \frac{i}{2\pi} \partial \bar{\partial} |w|^2 + \frac{-1}{\pi^2} \partial_z \bar{\partial}_z \partial_w \bar{\partial}_w G(e^{-\frac{1}{2}|z-w|^2}). \quad (37)$$

This completes the proof for the pair correlation case $n = 2$. (Notice that the formula has the same form in all dimensions.)

The proof for general n is similar. We again write $f^N = |f^N|u^N$ and expand the integrand in (23):

$$\begin{aligned} & \log |c \cdot f^N(\frac{z^1}{\sqrt{N}})| \log |c \cdot f^N(\frac{z^2}{\sqrt{N}})| \cdots \log |c \cdot f^N(\frac{z^n}{\sqrt{N}})| \\ &= \log |f^N(\frac{z^1}{\sqrt{N}})| \log |f^N(\frac{z^2}{\sqrt{N}})| \cdots \log |f^N(\frac{z^n}{\sqrt{N}})| \\ &+ \log |f^N(\frac{z^1}{\sqrt{N}})| \log |f^N(\frac{z^2}{\sqrt{N}})| \cdots \log |f^N(\frac{z^{n-1}}{\sqrt{N}})| \log |c \cdot u^N(\frac{z^n}{\sqrt{N}})| \\ &+ \cdots \\ &+ \log |c \cdot u^N(\frac{z^1}{\sqrt{N}})| \log |c \cdot u^N(\frac{z^2}{\sqrt{N}})| \cdots \log |c \cdot u^N(\frac{z^n}{\sqrt{N}})|. \end{aligned}$$

We denote the terms resulting from this expansion by E_1, \dots, E_{2^n} , respectively. As before, the first term converges to the normalized Euclidean “ n -fold” Kähler form:

$$E_1 = \frac{i}{2\pi} \partial \bar{\partial} |z^1|^2 \wedge \dots \wedge \frac{i}{2\pi} \partial \bar{\partial} |z^n|^2 + O\left(\frac{1}{N}\right).$$

The E_{2^n} term is obtained from the function

$$G_n^N(x^1, x^2, \dots, x^n) := \int_{\mathbb{C}^{d_N}} \log |c \cdot x^1| \log |c \cdot x^2| \dots \log |c \cdot x^n| \frac{e^{-|c|^2}}{\pi^{d_N}} dc, \quad (38)$$

$x^1, x^2, \dots, x^n \in \mathbb{C}^{d_N}$. Precisely, we substitute

$$x^j = u^N\left(\frac{z^j}{\sqrt{N}}\right) \quad (39)$$

in (38) and apply the operator $\left(\frac{i}{\pi}\right)^n \partial_{z^1} \bar{\partial}_{z^1} \dots \partial_{z^n} \bar{\partial}_{z^n}$. As above, we define a special Hermitian orthonormal basis $\{e_1, \dots, e_n\}$ for the n -dimensional complex subspace spanned by $\{x_1, \dots, x_n\}$. We put:

$$\begin{aligned} x^1 &= e_1 \\ x^2 &= \xi_{21}e_1 + \xi_{22}e_2 & \xi_{22} &= \sqrt{1 - |\xi_{21}|^2} \\ &\vdots \\ x^n &= \xi_{n1}e_1 + \dots + \xi_{nn}e_n & \xi_{nn} &= \sqrt{1 - \sum_{j \leq n-1} |\xi_{nj}|^2}. \end{aligned}$$

Such a basis exists because we can always multiply e_j by a phase $e^{i\theta}$ so that the last component ξ_{jj} is positive real. We complete $\{e_j\}$ to a basis of \mathbb{C}^{d_N} , and we now let c_j denote coordinates relative to this basis. As above, we rewrite the Gaussian integral in these coordinates. After integrating out the variables $\{c_{n+1}, \dots, c_{d_N}\}$, (38) simplifies to the n -dimensional complex Gaussian integral

$$\begin{aligned} G_n^N(x^1, \dots, x^n) &= G_n(\xi_{21}, \xi_{22}, \dots, \xi_{nn}) \\ &= \frac{1}{\pi^n} \int_{\mathbb{C}^n} e^{-|c|^2} \log |c_1| \log |c_1 \xi_{21} + c_2 \xi_{22}| \dots \log |c_1 \xi_{n1} + \dots c_n \xi_{nn}| dc. \end{aligned} \quad (40)$$

Note that the variables ξ_{jk} depend on N ; we write $\xi_{jk} = \xi_{jk}^N$ when we need to indicate this dependence.

To prove universality, we observe that the ξ_{jk} are universal algebraic functions of the inner products $\langle x^a, x^b \rangle$. Indeed,

$$\xi_{j1} \bar{\xi}_{k1} + \dots + \xi_{jk} \bar{\xi}_{kk} = \langle x^j, x^k \rangle, \quad 1 \leq k \leq j \leq n, \quad (41)$$

where we set $\xi_{11} = 1$. These algebraic functions are obtained by induction (lexicographically) using (41). (The triangular matrix (ξ_{jk}) is just the inverse of the matrix describing the Gram-Schmidt process.)

By (39), it follows that the ξ_{jk}^N are universal algebraic functions of the variables

$$\begin{aligned} \left\langle u^N\left(\frac{z^j}{\sqrt{N}}\right), u^N\left(\frac{z^k}{\sqrt{N}}\right) \right\rangle &= \frac{\Pi_N\left(\frac{z^j}{\sqrt{N}}, \frac{z^k}{\sqrt{N}}\right)}{\Pi_N\left(\frac{z^j}{\sqrt{N}}, \frac{z^j}{\sqrt{N}}\right)^{1/2} \Pi_N\left(\frac{z^k}{\sqrt{N}}, \frac{z^k}{\sqrt{N}}\right)^{1/2}} \\ &= e^{i\Im(z^j \cdot \bar{z}^k) - \frac{1}{2}|z^j - z^k|^2} + O\left(\frac{1}{\sqrt{N}}\right). \end{aligned}$$

We note here that

$$\begin{aligned} |x^1 \wedge \cdots \wedge x^n|^2 &= \det(\langle x^j, x^k \rangle) \rightarrow \det \left(e^{i\Im(z^j \cdot \bar{z}^k)} - \frac{1}{2}|z^j - z^k|^2 \right) \\ &= e^{-\sum |z^j|^2} \det \left(e^{z^j \cdot \bar{z}^k} \right). \end{aligned} \tag{42}$$

When the z_j are distinct (i.e., $(z^1, \dots, z^n) \in \mathcal{G}_n^m$), the limit determinant in (42) is nonzero (see [BSZ]) and thus $\xi_{jk}^N = \xi_{jk}^\infty + O(\frac{1}{\sqrt{N}})$, where the ξ_{jk}^∞ are universal real-analytic functions of $z \in \mathcal{G}_n^m$. We conclude that the E_{2^n} term converges to a universal current:

$$E_{2^n} = \left(\frac{i}{\pi} \right)^n \partial_{z^1} \bar{\partial}_{z^1} \cdots \partial_{z^n} \bar{\partial}_{z^n} G_n(\xi_{21}^\infty, \dots, \xi_{kk}^\infty) + O(\frac{1}{\sqrt{N}}).$$

Consider now a general term E_a . Suppose without loss of generality that E_a comes from

$$\log |c \cdot u^N(\frac{z^1}{\sqrt{N}})| \cdots \log |c \cdot u^N(\frac{z^k}{\sqrt{N}})| \log |f^N(\frac{z^{k+1}}{\sqrt{N}})| \cdots \log |f^N(\frac{z^n}{\sqrt{N}})|.$$

As above we obtain

$$\begin{aligned} E_a &= \left(\frac{i}{\pi} \right)^k \partial_{z^1} \bar{\partial}_{z^1} \cdots \partial_{z^k} \bar{\partial}_{z^k} G_k(\xi_{21}^\infty, \dots, \xi_{kk}^\infty) \wedge \frac{i}{2\pi} \partial \bar{\partial} |z^{k+1}|^2 \wedge \\ &\quad \cdots \wedge \frac{i}{2\pi} \partial \bar{\partial} |z^n|^2 + O(\frac{1}{\sqrt{N}}). \end{aligned}$$

Hence this term also approaches a universal current. (As in the pair correlation case, terms with only one u^N vanish.) \square

4. Explicit Formulae

We now calculate explicitly the limit pair correlation measures $\tilde{K}_2^\infty(z, w)$.

4.1. Preliminaries. The first step is to compute $\Delta G(e^{-\frac{1}{2}r^2})$, where Δ is the Euclidean Laplacian on \mathbb{C}^m and $r = |\zeta|$ ($\zeta \in \mathbb{C}^m$). To begin this computation, we write $a_j = r_j e^{i\varphi_j}$ and then rewrite (33)–(34) as

$$G(\cos \theta) = \frac{2}{\pi} \int_0^\infty \int_0^\infty \int_0^{2\pi} r_1 r_2 e^{-(r_1^2 + r_2^2)} \log r_1 \log |r_1 \cos \theta + r_2 e^{i\varphi} \sin \theta| d\varphi dr_1 dr_2. \tag{43}$$

We now evaluate the inner integral by Jensen’s formula, which gives

$$\int_0^{2\pi} \log |r_1 \cos \theta + r_2 \sin \theta e^{i\varphi}| d\varphi = \begin{cases} 2\pi \log(r_1 \cos \theta) & \text{for } r_2 \sin \theta \leq r_1 \cos \theta \\ 2\pi \log(r_2 \sin \theta) & \text{for } r_2 \sin \theta \geq r_1 \cos \theta \end{cases}. \tag{44}$$

Hence

$$G(\cos \theta) = 4 \int_0^\infty \int_0^\infty r_1 r_2 e^{-(r_1^2+r_2^2)} \log r_1 \log \max(r_1 \cos \theta, r_2 \sin \theta) dr_1 dr_2. \quad (45)$$

Now change variables again with $r_1 = \rho \cos \varphi$, $r_2 = \rho \sin \varphi$ to get

$$G(\cos \theta) = 4 \int_0^\infty \int_0^{\pi/2} \rho^3 e^{-\rho^2} \log(\rho \cos \varphi) \log \max(\rho \cos \varphi \cos \theta, \rho \sin \varphi \sin \theta) \cdot \cos \varphi \sin \varphi d\varphi d\rho. \quad (46)$$

Since

$$\log \max(\rho \cos \varphi \cos \theta, \rho \sin \varphi \sin \theta) = \log(\rho \cos \varphi \cos \theta) + \log^+(\tan \varphi \tan \theta),$$

we can write $G = G_1 + G_2$, where

$$G_1(\cos \theta) = 4 \int_0^\infty \int_0^{\pi/2} \rho^3 e^{-\rho^2} \log(\rho \cos \varphi) \log(\rho \cos \varphi \cos \theta) \cos \varphi \sin \varphi d\varphi d\rho, \quad (47)$$

$$G_2(\cos \theta) = 4 \int_0^\infty \int_{\pi/2-\theta}^{\pi/2} \rho^3 e^{-\rho^2} \log(\rho \cos \varphi) \log(\tan \varphi \tan \theta) \cos \varphi \sin \varphi d\varphi d\rho. \quad (48)$$

From (47), $G_1(\cos \theta) = C_1 + C_2 \log \cos \theta$ and thus

$$G_1(e^{-\frac{1}{2}r^2}) = C_1 - \frac{1}{2}C_2 r^2,$$

so that

$$\Delta G_1(e^{-\frac{1}{2}r^2}) = \left(\frac{d^2}{dr^2} + \frac{2m-1}{r} \frac{d}{dr} \right) (C_1 - \frac{1}{2}C_2 r^2) = -2mC_2. \quad (49)$$

We now evaluate $\Delta G_2(e^{-\frac{1}{2}r^2})$. Since the integrand in (48) vanishes when $\varphi = \pi/2 - \theta$, we have

$$\frac{d}{dr} G_2(\cos \theta) = 4 \left(\frac{d}{dr} \log \tan \theta \right) \int_0^\infty \int_{\pi/2-\theta}^{\pi/2} \rho^3 e^{-\rho^2} \log(\rho \cos \varphi) \cos \varphi \sin \varphi d\varphi d\rho.$$

Substituting $\tan^2 \theta = e^{r^2} - 1$, we have

$$\frac{d}{dr} \log \tan \theta = \frac{r}{1 - e^{-r^2}}.$$

Thus

$$\frac{d}{dr} G_2(e^{-\frac{1}{2}r^2}) = \frac{4r}{1 - e^{-r^2}} (I_1 + I_2),$$

where

$$I_1 = \int_0^\infty \int_{\pi/2-\theta}^{\pi/2} \rho^3 e^{-\rho^2} (\log \rho) \cos \varphi \sin \varphi d\varphi d\rho = C \sin^2 \theta = C(1 - e^{-r^2}),$$

$$I_2 = \int_0^\infty \int_{\pi/2-\theta}^{\pi/2} \rho^3 e^{-\rho^2} (\log \cos \varphi) \cos \varphi \sin \varphi d\varphi d\rho.$$

We compute

$$\begin{aligned} I_2 &= \frac{1}{2} \int_{\pi/2-\theta}^{\pi/2} (\log \cos \varphi) \cos \varphi \sin \varphi d\varphi = \frac{1}{2} \int_0^{\sin \theta} t \log t dt \\ &= \frac{1}{8} (\sin^2 \theta \log \sin^2 \theta - \sin^2 \theta) = \frac{1}{8} (1 - e^{-r^2}) [\log(1 - e^{-r^2}) - 1]. \end{aligned}$$

Thus

$$\frac{d}{dr} G_2(e^{-\frac{1}{2}r^2}) = \frac{r}{2} \log(1 - e^{-r^2}) + C'r. \tag{50}$$

Hence by (49) and (50),

$$\begin{aligned} \Delta G(e^{-\frac{1}{2}r^2}) &= -2mC_2 + \left(\frac{d}{dr} + \frac{2m-1}{r} \right) \left(\frac{r}{2} \log(1 - e^{-r^2}) + C'r \right) \\ &= m \log(1 - e^{-r^2}) + \frac{r^2}{e^{r^2} - 1} + C''. \end{aligned} \tag{51}$$

4.2. *Pair correlation in dimension 1.* In dimension one, the pair correlation form is the same as the pair correlation measure. We first give our universal formula in the one-dimensional case. Our formula agrees with that of Hannay [Ha] for SU(2) polynomials.

Theorem 4.1. *Suppose $\dim M = 1$. Then*

$$\begin{aligned} \mathbf{K}_2^N \left(\frac{z}{\sqrt{N}}, \frac{w}{\sqrt{N}} \right) &\rightarrow \mathbf{K}_2^\infty(z, w) = \left[\pi \delta_0(z - w) + H\left(\frac{1}{2}|z - w|^2\right) \right] \\ &\quad \cdot \frac{i}{2\pi} \partial \bar{\partial} |z|^2 \wedge \frac{i}{2\pi} \partial \bar{\partial} |w|^2, \end{aligned}$$

where

$$H(t) = \frac{(\sinh^2 t + t^2) \cosh t - 2t \sinh t}{\sinh^3 t} = t - \frac{2}{9}t^3 + \frac{2}{45}t^5 + O(t^7).$$

Proof. Making the change of variables $\zeta = z - w$, we have by (37),

$$\begin{aligned} \mathbf{E} \left(\widehat{Z}^N(z) \otimes \widehat{Z}^N(w) \right) &\rightarrow \frac{i}{2\pi} \partial \bar{\partial} |z|^2 \wedge \frac{i}{2\pi} \partial \bar{\partial} |w|^2 - \frac{1}{\pi^2} \partial_z \bar{\partial}_z \partial_w \bar{\partial}_w G(e^{-\frac{1}{2}|z-w|^2}) \\ &= \left[1 + 4 \frac{\partial^2}{\partial z \bar{\partial} z} \frac{\partial^2}{\partial w \bar{\partial} w} G(e^{-\frac{1}{2}|z-w|^2}) \right] \frac{i}{2\pi} \partial \bar{\partial} |z|^2 \wedge \frac{i}{2\pi} \partial \bar{\partial} |w|^2 \\ &= \left[1 + 4 \left(\frac{\partial^2}{\partial \zeta \bar{\partial} \zeta} \right)^2 G(e^{-\frac{1}{2}|\zeta|^2}) \right] \frac{i}{2\pi} \partial \bar{\partial} |z|^2 \wedge \frac{i}{2\pi} \partial \bar{\partial} |w|^2 \\ &= \left[1 + \frac{1}{4} \Delta^2 G(e^{-\frac{1}{2}r^2}) \right] \frac{i}{2\pi} \partial \bar{\partial} |z|^2 \wedge \frac{i}{2\pi} \partial \bar{\partial} |w|^2. \end{aligned}$$

By (51) with $m = 1$, we have

$$\begin{aligned} \Delta^2 G(e^{-\frac{1}{2}r^2}) &= \left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} \right) \left[\log(1 - e^{-r^2}) + \frac{r^2}{e^{r^2} - 1} \right] \\ &= 4\pi\delta_0 + \frac{8(e^{r^2} - 1)^2 - 16r^2 e^{r^2} (e^{r^2} - 1) + 4r^4 e^{r^2} (e^{r^2} + 1)}{(e^{r^2} - 1)^3}. \end{aligned}$$

Finally,

$$\begin{aligned} \left[1 + \frac{1}{4} \Delta^2 G(e^{-\frac{1}{2}r^2}) \right] &= \pi\delta_0 + \frac{(e^{r^2} + 1)(e^{r^2} - 1)^2 - 4r^2 e^{r^2} (e^{r^2} - 1) + r^4 e^{r^2} (e^{r^2} + 1)}{(e^{r^2} - 1)^3} \\ &= \pi\delta_0 + \frac{(\sinh^2 \frac{1}{2}r^2 + \frac{1}{4}r^4) \cosh \frac{1}{2}r^2 - r^2 \sinh \frac{1}{2}r^2}{\sinh^3 \frac{1}{2}r^2}. \quad \square \end{aligned}$$

4.3. *Pair correlation in higher dimensions.* The limit pair correlation measure is given by

$$\begin{aligned} \tilde{K}_2^\infty(z, w) &= \lim_{N \rightarrow \infty} N^{2(m-1)} \tilde{K}_2^N\left(\frac{z}{\sqrt{N}}, \frac{w}{\sqrt{N}}\right) \\ &= \mathbf{K}_2^\infty(z, w) \wedge \frac{1}{(m-1)!} \left(\frac{i}{2} \partial \bar{\partial} |z|^2\right)^{m-1} \wedge \frac{1}{(m-1)!} \left(\frac{i}{2} \partial \bar{\partial} |w|^2\right)^{m-1}. \end{aligned}$$

(The scaling $N^{2(m-1)}$ comes from the fact that $N\omega(\frac{z}{\sqrt{N}}) = N(\tau_{\sqrt{N}})_*\omega \rightarrow \frac{i}{2} \partial \bar{\partial} |z|^2$.)

We now compute \tilde{K}_2^∞ for the case of a manifold of general dimension $m > 1$. It is convenient to express this measure in terms of the expected density of zeros

$$\tilde{K}_1^\infty(z) = \lim_{N \rightarrow \infty} N^{m-1} \tilde{K}_1^N\left(\frac{z}{\sqrt{N}}\right) = \frac{m}{\pi} dV_{\mathbb{C}^m} = \frac{1}{\pi(m-1)!} \left(\frac{i}{2} \partial \bar{\partial} |z|^2\right)^m. \quad (52)$$

We have the following explicit universal formula for the limit pair correlation measure. In particular, it gives the scaling limit pair correlation for the zeros of $SU(m + 1)$ -polynomials.

Theorem 4.2. *Suppose $\dim M = m > 1$. Then*

$$\tilde{K}_2^\infty(z, w) = \left[\gamma_m\left(\frac{1}{2}|z - w|^2\right) \right] \tilde{K}_1^\infty(z) \wedge \tilde{K}_1^\infty(w),$$

where

$$\begin{aligned} \gamma_m(t) &= \frac{\left[\frac{1}{2}(m^2 + m) \sinh^2 t + t^2 \right] \cosh t - (m + 1)t \sinh t}{m^2 \sinh^3 t} + \frac{m - 1}{2m} \\ &= \frac{(m - 1)}{2m} t^{-1} + \frac{m - 1}{2m} + \frac{(m + 2)(m + 1)}{6m^2} t \\ &\quad - \frac{(m + 4)(m + 3)}{90m^2} t^3 + \frac{(m + 6)(m + 5)}{945m^2} t^5 + O(t^7). \end{aligned}$$

Proof. By (37) and (51), again writing $\zeta = z - w$ (except this time $\zeta \in \mathbb{C}^m$),

$$\begin{aligned} \tilde{K}_2^\infty(z, w) &= \left[1 + \frac{4}{m^2} \sum_{j,k=1}^m \frac{\partial^2}{\partial z_j \partial \bar{z}_j} \frac{\partial^2}{\partial w_k \partial \bar{w}_k} G(e^{-\frac{1}{2}|z-w|^2}) \right] \tilde{K}_1^\infty(z) \wedge \tilde{K}_1^\infty(w) \\ &= \left[1 + \frac{1}{4m^2} \Delta_\zeta^2 G(e^{-\frac{1}{2}|\zeta|^2}) \right] \tilde{K}_1^\infty(z) \wedge \tilde{K}_1^\infty(w) \\ &= \left[1 + \frac{1}{4m^2} \left(\frac{d^2}{dr^2} + \frac{2m-1}{r} \frac{d}{dr} \right) \left(m \log(1 - e^{-r^2}) + \frac{r^2}{e^{r^2} - 1} \right) \right] \tilde{K}_1^\infty(z) \wedge \tilde{K}_1^\infty(w). \end{aligned} \quad (53)$$

Computing the Laplacian in (53) leads to the stated formula. \square

Note that if we substitute $m = 1$ in the expression for $\gamma_m(t)$, we obtain Hannay's function $H(t)$. However for the case $m > 1$, the limit measure is absolutely continuous on $\mathbb{C}^m \times \mathbb{C}^m$, whereas in the one-dimensional case, there is a self-correlation delta measure.

Acknowledgement. The first draft of this paper was completed while the third author was visiting the Erwin Schrödinger Institute in July 1998. He wishes to thank that institution for its hospitality and financial support.

References

- [BD] Bleher, P. and Di, X.: Correlations between zeros of a random polynomial. *J. Stat. Phys.* **88**, 269–305 (1997)
- [BSZ] Bleher, P., Shiffman, B. and Zelditch, S.: Universality and scaling of correlations between zeros on complex manifolds. *Invent. Math.* to appear
- [BBL] Bogomolny, E., Bohigas, O. and Leboeuf, P.: Quantum chaotic dynamics and random polynomials. *J. Stat. Phys.* **85**, 639–679 (1996)
- [D] Deift, P.: *Orthogonal polynomials and random matrices: A Riemann–Hilbert approach*. Courant Lecture Notes in Mathematics, **3**, New York: New York University, Courant Institute of Mathematical Sciences, 1999
- [GH] Griffiths, P. and Harris, J.: *Principles of Algebraic Geometry*. New York: Wiley-Interscience, 1978
- [Ha] Hannay, J.H.: Chaotic analytic zero points: Exact statistics for those of a random spin state. *J. Phys. A: Math. Gen.* **29**, 101–105 (1996)
- [SZ] Shiffman, B. and Zelditch, S.: Distribution of zeros of random and quantum chaotic sections of positive line bundles. *Commun. Math. Phys.* **200**, 661–683 (1999)
- [Ti] Tian, G.: On a set of polarized Kähler metrics on algebraic manifolds. *J. Diff. Geometry* **32**, 99–130 (1990)
- [Ze] Zelditch, S.: Szegő kernels and a theorem of Tian. *Int. Math. Res. Notices* **6**, 317–331 (1998)

Communicated by P. Sarnak